



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

vanishes for $u \neq 0$. Thus (7) has at most one positive solution, and similarly at most one negative solution.

Finally let $\alpha = 0$. Equations (4), (3) reduce to

$$3 \sin \beta l = \beta(1 + \cos \beta l),$$

$$\beta \sin \beta l = 3(1 - \cos \beta l);$$

and this pair of equations is equivalent to the single equation

$$3 \sin \frac{1}{2}\beta l = \beta \cos \frac{1}{2}\beta l.$$

Since it is clear that no value which makes $\cos \frac{1}{2}\beta l$ vanish can be a solution, the only admissible values of β are those for which

$$\tan \frac{1}{2}\beta l = \frac{1}{3}\beta, \quad (8)$$

which may be written

$$\tan u - ku = 0, \quad (9)$$

where

$$u = \frac{1}{2}\beta l, \quad k = \frac{2}{3l}.$$

This is a type of equation which arises in many problems of analysis. Inspection of the graphs of $\tan u$ and ku leads at once to the well-known fact, which can readily be proved by rigorous analytic methods, that (8) has an infinite number of solutions, approximating more and more closely, as they increase numerically, to odd multiples of $\pi/2$.

We condense all these results into the statement: The function x^n , where n is a constant whose real part is positive, satisfies the proposed functional equation for the following values of n and no others:

(a) The real values $n = 1, 2, 3$.

(b) An infinite set of values of the form $n = 2 + \beta i$, where β is a solution¹ of (8).

Any of the second set of solutions leads to real solutions of the type considered by Professor Bennett. In fact, since

$$x^n = x^{a+2} = x^{a+2+\beta i} = x^{a+2} \cos(\beta \log x) + ix^{a+2} \sin(\beta \log x),$$

for any n yielding a complex solution we have the two real solutions $x^{a+2} \cos(\beta \log x)$ and $x^{a+2} \sin(\beta \log x)$, whence also the more general solution $x^{a+2} \sin(\beta \log x - c)$, which is a linear combination of them. Equations (3), (4) are equivalent to Professor Bennett's pair of equations for a, b with $a = \alpha + 2$, $b = \beta$. We have thus shown that Professor Bennett's formula gives a solution only (except for the trivial cases $b = 0$, $a = 1, 2, 3$) when $a = 2$, and that in this instance an infinite number of values of b will be possible. It is obvious that much more general solutions can be constructed, of the form $x^2 F(x)$, where

$$F(x) = \Sigma[A_p \cos(\beta_p \log x) + B_p \sin(\beta_p \log x)],$$

proper precautions being taken to insure the validity of the series.

The importance of these new solutions lies in the following facts. The function $x^2 \sin(\beta \log x - c)$ possesses a continuous first derivative even at the origin. In previous editorial comment, it was stated that "it seems reasonable to adopt as a goal with reference to this question the proof that if the equation holds for some range of the variable h , the function $F(x)$ can only be a polynomial of degree ≤ 3 , under restrictions as light as possible—*e.g.*, that $F(x)$ should be continuous and possess a stated number (as small as it can be made) of derivatives." The statement has now been proved, with the number of derivatives as great as *six*, by Professor Gillespie. Professor Bennett's example, together with the present comments, shows that the theorem is not true with the number of derivatives as small as *one*. Can the gap between these two results be bridged?

DISCUSSIONS.

In a previous number of the MONTHLY (1920, 53) Mr. W. F. Cheney gave an instance of a geometric proof of the law of tangents in trigonometry. In the first discussion below Professor Lovitt shows how a number of such proofs may be obtained. The two papers contain references to a number of other similar proofs.

¹ The smallest positive value of β is approximately 12.94.

